


temperature (3-5°C). Filamentous fungi were maintained on Czapek Malt slants at refrigeration temperature (3-5°C), until needed for the assay.

In the Claims:

Please cancel claims 1-32.

Please add new claims 33-43 as follows.

~~33.~~ (New) An oleanane triterpenoid oligoglycoside isolated from *Aegiceras corniculatum* and having the formula I:



wherein R₁ is selected from the group consisting of -OH, -OAc and =O.

~~34.~~ (New) The isolated oleanane triterpenoid oligoglycoside compound according to claim 1 wherein R₁ has a α or β configuration.

²
~~34.~~ (Amended) The isolated oleanane triterpenoid oligoglycoside compound according to claim ¹~~33~~ wherein R₁ has a α or β configuration.

³
~~35.~~ (Amended) The isolated oleanane triterpenoid oligoglycoside compound according to claim ¹~~33~~ wherein the polysaccharide carbohydrate portion has a monosaccharide having a D or L rotation and the intersugar linkages have a α or β configuration.

⁴
~~36.~~ (Amended) The isolated oleanane triterpenoid oligoglycoside compound according to claim ¹~~33~~ wherein the carbohydrate portion linked to the aglycone portion at position 3 has a β configuration.

⁵
~~37.~~ (Amended) The isolated oleanane triterpenoid oligoglycoside compound according to claim ¹~~33~~ wherein R₁ is -OH.

⁶
~~38.~~ (Amended) The isolated oleanane triterpenoid oligoglycoside compound according to claim ¹~~33~~ wherein the C₁₃ - C₂₈ oxydo ring has a β configuration.

⁷
~~39.~~ (Amended) The isolated oleanane triterpenoid oligoglycoside compound according to claim ¹~~33~~ wherein the compound inhibits growth of fungi selected from *candida* sp. and *Cryptococcus neoformans*.

⁸
~~40~~. (Amended) The isolated oleanane triterpenoid oligoglycoside compound according to claim ¹~~33~~ wherein the compound inhibits growth of *Microsporin gypseum*, *Trichophyton mentagrophytes*, *T. tonsurans* and *T. violaceum*.

⁹
~~41~~. (New) | The isolated oleanane triterpenoid oligoglycoside compound according to claim ³³ ~~1~~ wherein the compound inhibits growth of *Aspergillus fresenii* and *A. niger*.

¹⁰
~~42~~. (New) | The isolated oleanane triterpenoid oligoglycoside compound according to claim ³⁵ ~~1~~ wherein the compound is used for treating candidiasis, cryptococcosis, epidermal infections and systemic infections.

¹¹
~~43~~. (New) | The isolated oleanane triterpenoid oligoglycoside compound according to claim ³⁵ ~~1~~ wherein the compound is used as a food preservative.
